



Rancho Los Amigos

Post-Polio

Support Group

Newsletter - August 2005

Editor's Note: The following summary was part of the information presented at the Ninth International Conference on Post-Polio Health and Ventilator-Assisted Living, held in St. Louis, Missouri USA from June 2 to 4, 2005. It is reprinted here with the author's permission.

Summary of Anesthesia Issues for Post-Polio Patients

By Selma Harrison Calmes MD
Clinical Professor of Anesthesiology, UCLA

Polio results in wide-spread neural changes, not just destruction of the spinal cord anterior horn (motor nerve) cells, and these changes can get worse as patients age. These anatomic changes affect many aspects of anesthesia care. No study of polio patients having anesthesia has been done. These recommendations are based on extensive review of the current literature and clinical experience with these patients. They may need to be adjusted for a particular patient.

1. Post-polio patients are nearly always very sensitive to sedative meds, and emergence can be prolonged. This is probably due to central neuronal changes, especially in the Reticular Activating System, from the original disease.

2. Non-depolarizing muscle relaxants cause a greater degree of block for a longer period of time in post-polio patients. The current recommendation is to start with half the usual dose of whatever you're using, adding more as needed. This is because the poliovirus actually lived at the neuromuscular junctions during the original disease, and there are extensive anatomic changes there, even in seem-

(Continued on page 2)

Unless otherwise stated, the articles in this newsletter may be reprinted provided they are reproduced in their entirety, and the author, the original source, and the Rancho Los Amigos Post-Polio Support Group are acknowledged in full. **Copyrighted articles may not be used without prior approval, and no article may be edited.** The information presented at our meetings and/or contained in this newsletter is solely for information. It is not to be taken as an endorsement of any product, medication, or individual.

(Continued from page 1)

ingly normal muscles, which make for greater sensitivity to relaxants. Also, many patients have a significant decrease in total muscle mass. Neuromuscular monitoring intraop helps prevent overdose of muscle relaxants. Overdose has been a frequent problem.

3. Succinylcholine often causes severe, generalized muscle pain postop. It is useful if this can be avoided, if possible.

4. Postop pain is often a significant issue. The anatomic changes from the original disease can affect pain pathways due to “spill-over” of the inflammatory response. Spinal cord “wind-up” of pain signals seems to occur. Proactive, multimodal postop pain control (local anesthesia at the incision plus PCA, etc) helps.

5. The autonomic nervous system is often dysfunctional, again due to anatomic changes from the original disease (the inflammation and scarring in the anterior horn “spills over” to the intermediolateral column, where sympathetic nerves travel). This can cause gastro-esophageal reflux, tachyarrhythmias and, sometimes, difficulty maintaining BP when anesthetics are given.

6. Patients who use ventilators often have worsening of ventilatory function postop, and some patients who did not need assisted ventilation have had to go onto a ventilator (including long-term use) postop. It's useful to get at least a VC preop, and full pulmonary function studies may be helpful. One group that should all have preop PFTs is those who were in iron lungs. The marker for real difficulty is thought to be a VC < 1.0 liter. Such a patient needs good pulmonary preparation preop and a plan for postop ventilatory support. Another ventilation risk is obstructive sleep apnea in the postop period. Many post-polios are turning out to have significant sleep apnea due to new weakness in their upper airway muscles as they age.

7. Laryngeal and swallowing problems due to muscle weakness are being recognized more often. Many patients have at least one paralyzed cord, and several cases of bilateral cord paralysis have occurred postop, after intubation or upper extremity blocks. ENT evaluation of the upper airway in suspicious patients would be useful.

8. Positioning can be difficult due to body asymmetry. Affected limbs are

(Continued on page 3)

(Continued from page 2)

osteopenic and can be easily fractured during positioning for surgery. There seems to be greater risk for peripheral nerve damage (includes brachial plexus) during long cases, probably because nerves are not normal and also because peripheral nerves may be unprotected by the usual muscle mass or tendons.

⌘ ⌘ ⌘ ⌘ ⌘